

AN EASY CONSTRUCTION OF G_2

N. J. WILDBERGER

ABSTRACT. We show how to construct the simple exceptional Lie algebra of type G_2 by explicitly constructing its 7 dimensional representation. No knowledge of Lie theory is required, and all relevant coefficients can be remembered by use of the simple mnemonic $\beta\alpha\beta\beta\alpha\beta$.

1. INTRODUCTION

The classification of simple Lie algebras goes back to W. Killing [9] and E. Cartan [2]. There are four families of ‘classical’ Lie algebras of type A, B, C, D corresponding to the special linear, odd orthogonal, symplectic and even orthogonal algebras, and five exceptional Lie algebras of types G_2, F_4, E_6, E_7 and E_8 of dimensions 14, 56, 78, 133 and 248 respectively, which we denote by $\mathfrak{g}_2, \mathfrak{f}_4$ etc. The construction of these exceptional Lie algebras is subtle and has been treated by several authors beginning with Cartan. A general but cumbersome construction valid for all simple Lie algebras was given by Harish-Chandra [5], while for the exceptional Lie algebras Tits [13] has given a uniform but perhaps complicated treatment. In principle the Serre relations allow us to write down structure constants for any simple Lie algebra from the Dynkin diagram, but this is somewhat abstract. In many concrete applications one desires to know the explicit structure constants with respect to a convenient basis.

Jacobson [7] and [8] constructs \mathfrak{g}_2 as the Lie algebra of derivations of a certain non-associative Cayley algebra. Adams [1] shows that the corresponding Lie group G_2 is the subgroup of $Spin(7)$ fixing a point in S^7 . Humphreys [6] shows how to explicitly write down basis elements of \mathfrak{g}_2 in a somewhat ad hoc fashion, from knowledge of the simple Lie algebra of type B_3 . Fulton and Harris [4] devote most of a chapter to a construction using detailed facts from the representation theory of sl_2 and sl_3 .

Our purpose here is to give a quick and easy construction of \mathfrak{g}_2 , requiring *no knowledge of Lie theory*. Indeed the entire construction is given in this Introduction; the rest of the paper contains a discussion of connections with Lie theory, some relatively simple mnemonics for remembering the relevant graphs, a sketch of the proof of the main Theorem, and an explicit table of commutation relations.

Consider the directed graph of Figure 1, consisting of vertices around a hexagon together with its center, and certain directed edges (we adopt the lazy convention that an edge with two opposite arrows on it signifies two directed edges of opposite orientation.) We refer to this directed graph as the G_2 Hexagon, and note that in Figure 1 its directed edges are further labelled by integers which we call *weights* (the default weight being one). Each edge of the G_2 Hexagon has a direction given by one of the twelve vectors in Figure 2. We call these twelve vectors *roots* and note that each is an integral linear combination of the basis vectors α and β , in fact either with all nonnegative

coefficients, or all with non-positive coefficients. It will be useful to consider both Figures in the same plane and sharing a common origin.




Figure 1: The G_2 Hexagon




Figure 2: Roots of G_2

To each root γ we associate an operator X_γ on the vertices of the G_2 Hexagon, defined by the rule that it takes a vertex v to n times a vertex w precisely when there is an edge from v to w of direction γ and weight n . If we use the labelling of the vertices shown, then for example the operator X_α takes v_β to $v_{\beta\alpha}$ and takes $v_{\beta\alpha\beta\beta}$ to $v_{\beta\alpha\beta\beta\alpha}$, while it sends all other vertices to 0, while the operator $X_{\alpha+2\beta}$ takes v_ϕ to $2v_{\beta\alpha\beta}$, v_β to $-v_{\beta\alpha\beta\beta}$, $v_{\beta\alpha}$ to $-v_{\beta\alpha\beta\beta\alpha}$, and $v_{\beta\alpha\beta}$ to $v_{\beta\alpha\beta\beta\alpha\beta}$, while it sends the other vertices to 0. So we get twelve operators acting on the vector space V spanned by the vertices.

Let us agree that if X and Y are operators, XY denotes the composition operator $Y \circ X$ (do X first, then Y) and $[X, Y] = XY - YX$ denotes the bracket of operators. We now define some additional operators H_γ , one for each root γ , by the commutation relation $H_\gamma = [X_{-\gamma}, X_\gamma]$. It is not hard to verify that each of these H_γ is a *scalar* operator; it multiplies each vertex by a scalar, which in fact is the restriction of a linear function to the G_2 Hexagon, assumed to be centered at the origin.

Theorem 1.1. *The span of the operators $\{X_\gamma, H_\gamma \mid \gamma \text{ is a root}\}$ is closed under brackets and forms a 14-dimensional Lie algebra $\mathfrak{g} \simeq \mathfrak{g}_2$. A basis of \mathfrak{g}_2 consists of $\{X_\gamma \mid \gamma \text{ is a root}\} \cup \{H_\alpha, H_\beta\}$.*

With respect to the basis of vertices of the G_2 Hexagon, each of the operators X_γ , H_γ can be seen to have integer matrix entries in the set $\{-2, -1, 0, 1, 2\}$, with each row and column containing at most one nonzero entry. This implies that the structure equations of \mathfrak{g} can be read off easily from the G_2 Hexagon by simply observing how the operators compose. For example, $[X_\alpha, X_\beta] = -X_{\alpha+\beta}$, as the reader may quickly verify by checking the actions on a suitable vertex, such as ϕ .

The remainder of the paper shows that the entire procedure, including determination of the weights on the edges of the G_2 Hexagon, can be obtained by playing the ‘Mutation and Numbers Games’ on a particular graph. These games are of considerable independent interest, leading quickly to deep aspects of Lie theory. The Numbers Game was defined by Mozes [10], and studied by Proctor [11] and Eriksson [3], while the Mutation Game is in a precise sense dual to the Numbers Game and has its origins in the theory of reflection groups and root systems (see [16]). Ultimately the entire construction boils down to considering convex subsets of the chain $\beta < \alpha < \beta < \beta < \alpha < \beta$. Included is a full list of all commutation relations.

To agree with standard usage, we split the set of operators $\{X_\gamma \mid \gamma \text{ is a root}\}$ into operators X_γ, Y_γ for γ a positive root. These are then raising and lowering operators in the familiar terminology of the physics literature.

The construction presented here is in the same spirit as [14], in which we constructed the simply laced Lie algebras from their Dynkin diagrams (also without explicit use of Lie theory), and is related to two recent constructions of the irreducible representations of $sl(3)$ (different from Gelfand Tsetlin) called the *Diamond* and *Zed* Models in [15]. It is because of these connections that we describe the construction in terms of raising and lowering operators in ideals of posets associated to the Mutation and Numbers games.

2. CONNECTIONS WITH LIE THEORY

Consider the directed graph G_2 consisting of two vertices α and β with three edges from α to β and one edge from β to α . We may encapsulate this information as in Figure 3.




Figure 3: The directed graph G_2

We will now show that both Figure 1 and Figure 2 arise by playing two remarkable graph-theoretical games, called the Mutation Game and the Numbers Game, on the graph G_2 .

Given any directed graph Z , we may associate to it a distinguished collection $R = R(Z)$ of integer valued functions on the vertices of Z by playing the *Mutation Game*. The elements of R are called *roots*. A root is any function obtained by starting with a delta function— with value one at some vertex and zero elsewhere— and performing an arbitrary sequence of *mutations*, defined as follows. A mutation at a vertex z applied to a function f has the effect of leaving the values of f at all vertices except z unchanged, while the new value at z is obtained by negating the current value at z and adding a copy of the function values at each neighbour w of z , once for every directed edge from w to z . Since a mutation at z applied to the delta function at z results in a sign change, the set of roots is symmetric under change of sign.

The reader may now check that in our example the set $R(G_2)$ consists of the following *positive* roots together with their negatives.

$$R^+(G_2) = \{ \alpha, \beta, \alpha + \beta, \alpha + 2\beta, \alpha + 3\beta, 2\alpha + 3\beta \}.$$

As functions on G_2 , these are obtained from the delta functions α and β (usually referred to as *simple roots*), by mutations in the orders $\alpha \rightarrow \alpha + 3\beta \rightarrow 2\alpha + 3\beta$ and $\beta \rightarrow \alpha + \beta \rightarrow \alpha + 2\beta$. We now associate α and β to two vectors in the plane whose lengths are in the ratio $1 : \sqrt{3}$ respectively, making an angle of $5\pi/6$ as in Figure 2. The other roots, which are linear combinations of α and β , correspond to the other vectors in the same picture, and the mutations at the vertices α and β act on these vectors by reflections in the lines perpendicular to α and β respectively. Thus $R(G_2)$ is indeed a *root system* in the classical sense.

The *Numbers Game* is also played with integer valued functions on the vertices of a directed graph. Given such a function g and a vertex z , we will say that a *dual mutation* at z replaces the value of g at z with its negative, simultaneously adds a copy of $g(z)$ to each neighbour w of z , once for each directed edge from z to w , and leaves all other vertices unchanged. Of particular interest is playing the Numbers Game with the condition that dual mutations only can occur at vertices where the current value of the function is positive. The only example relevant for us here starts with a delta function at β , here denoted by recording the function values in the ordered pair $(0, 1)$. Proceeding with positive dual mutations, we get the sequence $(0, 1), (1, -1), (-1, 2), (1, -2), (-1, 1), (0, -1)$. If we record, using multiplicative notation, those positive function values at each stage in this dual mutation sequence, we get what we call the ‘change sequence’ $\beta\alpha\beta^2\alpha\beta$, which we write as $\beta\alpha\beta\beta\alpha\beta$. For

other graphs and other starting values, change sequences associated to such positive dual mutation sequences have remarkable combinatorial properties (see [13]).

Form a partially ordered multiset P by ordering the terms linearly by $\beta < \alpha < \beta < \beta < \alpha < \beta$. This poset (perhaps it ought to be called a *pomset*) is then a chain. Recall that a subset I of a poset is called an *ideal* if $x \in I$ and $y \leq x$ implies $y \in I$. We will consider the seven ideals of P to form the vertices of a graph in the plane whose edges have ‘directions’ corresponding to roots. More specifically we will connect an ideal v to an ideal u by an edge of *direction* γ precisely when $\sum_{x \in u} x - \sum_{x \in v} x = \gamma$. Thus for example there is an edge of direction α from $\beta\alpha\beta\beta$ to $\beta\alpha\beta\beta\alpha$, and an edge of direction $-\alpha - 2\beta$ from $\beta\alpha\beta\beta$ to β . This yields the G_2 Hexagon. It remains to prescribe the *weight* $w(e)$ of a directed edge e .

A subset $L \subseteq P$ is called a *layer* if it is convex in the usual sense. This means here only that L is a chain of some consecutively ordered elements of P , such as for example $\beta\beta\alpha\beta$ or $\alpha\beta\beta$. Such a layer L will be called a γ -*layer*, for some root $\gamma \in R^+$, if

$$\sum_{z \in L} z = \gamma.$$

We let \mathcal{L}_γ denote the set of γ -layers, which itself is partially ordered by setting $L_1 \leq L_2$ if $L_1 \subseteq I(L_2)$ where $I(S)$ is the ideal generated by a subset $S \subseteq P$, that is,

$$I(S) = \{z \mid z \leq w \text{ for some } w \in S\}$$

For example if $\gamma = \alpha + 2\beta$, then \mathcal{L}_γ contains four γ -layers; $L_0 = \beta\alpha\beta < L_1 = \alpha\beta\beta < L_2 = \beta\beta\alpha < L_3 = \beta\alpha\beta$. (Note that there is some possibility for confusion here as L_0 and L_3 are distinct layers despite the fact that as sequences they appear the same.) For each γ there is a unique minimal γ -layer which we will denote by L_m , and a unique maximal γ -layer which we will denote by L_M .

Given a γ -layer L , define the *parity* $\epsilon(L)$ (respectively *dual parity* $\tilde{\epsilon}(L)$) of L , by $(-1)^n$ where n is the number of $\alpha - \beta$ interchanges required in passing from L to the minimal γ -layer L_m (respectively maximal γ -layer L_M) by switching adjacent α and β 's. This is well-defined. Thus continuing the example above, $\epsilon(L_0) = 1$, $\epsilon(L_1) = -1$, $\epsilon(L_2) = -1$, and $\epsilon(L_3) = 1$, while the dual parities agree with the parities since L_m and L_M have the same form. A *ladder* of γ -layers will be a sequence $L_1 < L_2 < \dots < L_k$ of disjoint γ -layers such that $L_1 \cup \dots \cup L_i$ is a layer for all $i = 1, \dots, k$. Such a ladder has *size* k , and will be said to *start* at L_1 and *finish* at L_k .

For a γ -layer L let $s(L)$ be the maximum size of a ladder starting at L , and let $f(L)$ be the maximum size of a ladder finishing at L . Thus in the example above $s(L_0) = 2$, $s(L_1) = s(L_2) = s(L_3) = 1$, while $f(L_0) = f(L_1) = f(L_2) = 1$, $f(L_3) = 2$.

For a positive root γ and a γ -layer L , define the *weight* $w(L)$ and the *dual weight* $\tilde{w}(L)$ by

$$w(L) = \epsilon(L)s(L) \qquad \tilde{w}(L) = \tilde{\epsilon}(L)f(L).$$

Ideals in P correspond to vertices in the G_2 Hexagon, and γ -layers for $\gamma \in R$ to γ -edges, so we associate to a γ -edge e the weight $w(L)$ of the corresponding γ -layer. For a $-\gamma$ -edge of direction the negative of a positive root γ we associate the dual weight $\tilde{w}(L)$ of the corresponding γ -layer. The G_2 Hexagon is symmetrical about the origin, and this definition produces a pattern of weights on the G_2 Hexagon which is still symmetrical about the origin.

We now remind the reader about some elementary facts about the structure of a semisimple Lie algebra \mathfrak{g} . This is intended to motivate what follows, so is not strictly necessary for the construction itself. If R is the root system of \mathfrak{g} , then there exists a decomposition

$$\mathfrak{g} = \mathfrak{h} \oplus \sum_{\gamma \in R} \mathfrak{g}_{\gamma}$$

where \mathfrak{h} is a *Cartan subalgebra* (maximal commutative subalgebra of semisimple elements), and each \mathfrak{g}_{γ} is a one dimensional *root space* with the property that $[H, X_{\gamma}]$ is always a multiple of X_{γ} for $H \in \mathfrak{h}$ and $X_{\gamma} \in \mathfrak{g}_{\gamma}$.

Suppose one chooses, perhaps arbitrarily, a nonzero element X_{γ} from each root space \mathfrak{g}_{γ} . Then from general considerations (see for example Humphreys [H]) one knows that for $\gamma, \gamma' \in R$, $[X_{\gamma}, X_{\gamma'}] = 0$ unless $\gamma + \gamma'$ is also a root, in which case $[X_{\gamma}, X_{\gamma'}]$ is some multiple of $X_{\gamma+\gamma'}$, or unless $\gamma = -\gamma'$, in which case $[X_{\gamma}, X_{-\gamma}]$ is a known element of the Cartan subalgebra \mathfrak{h} . So the problem involved in describing \mathfrak{g} completely essentially involves two steps: to choose explicit basis elements X_{γ} in each root space \mathfrak{g}_{γ} and to determine the corresponding constants $n_{\gamma, \gamma'}$ in the structural equations

$$[X_{\gamma}, X_{\gamma'}] = n_{\gamma, \gamma'} X_{\gamma+\gamma'},$$

one for each pair of roots (γ, γ') whose sum is also a root. Then all other structure constants of the Lie algebra will be given. Somewhat surprisingly, the difficulty in this task lies not only in finding the absolute values of the $n_{\gamma, \gamma'}$; the signs can also be subtle to determine. See Samelson [12], Tits [13] for a discussion of this important point.

3. THE 7-DIMENSIONAL REPRESENTATION OF \mathfrak{g}_2

The Lie algebra \mathfrak{g}_2 will be realized by operators on the vector space spanned by the vertices of the G_2 Hexagon, which we have seen are labelled by the ideals in the poset P arising from playing the Numbers Game on a particular initial function. To abide by the usual conventions, we will actually associate to each positive root $\gamma \in R^+$ two operators, X_{γ} and Y_{γ} , corresponding to raising and lowering operators in the physics literature. Together with two scalar operators H_{α} and H_{β} , we get a total of fourteen operators on a seven dimensional space. All of these operators are entirely encoded in the graph and its weighted directed edges and we will then check that they form a Lie algebra of operators. This will be \mathfrak{g}_2 . We will show that if we augment our set of operators to a larger set of 18, still in the span of the original 14, then we obtain what we call a *bracket set*; the bracket of any two of them is a multiple of one in the set.

Let us define a complex vector space

$$V = \text{span}\{v_I \mid I \text{ is an ideal of } P\}.$$

For any layer $L \subseteq P$ define operators X_L, Y_L on V by

$$X_L(v_I) = \begin{cases} v_{I \cup L} & \text{if } I \cup L \text{ is an ideal and } I \cap L = \emptyset \\ 0 & \text{otherwise.} \end{cases}$$

$$Y_L(v_I) = \begin{cases} v_{I \setminus L} & \text{if } L \subseteq I \text{ and } I \setminus L \text{ is an ideal} \\ 0 & \text{otherwise.} \end{cases}$$

For $\gamma \in R^+$ define operators X_γ, Y_γ and H_γ on V by

$$X_\gamma = \sum_{L \in \mathcal{L}_\gamma} w(L)X_L \quad Y_\gamma = \sum_{L \in \mathcal{L}_\gamma} w(\tau(L))y_L$$

$$H_\gamma(v_I) = \begin{cases} kv_I & \text{if } k \text{ is the largest positive integer such that there exists a ladder of } \gamma\text{-layers} \\ & L_1 < \dots < L_k \subseteq I \text{ such that } I \setminus (L_1 \cup \dots \cup L_k) \text{ is an ideal} \\ -kv_I & \text{if } k \text{ is the largest positive integer such that there exists a ladder of } \gamma\text{-layers} \\ & I < L_1 < \dots < L_k \text{ such that } I \cup (L_1 \cup \dots \cup L_k) \text{ is an ideal and} \\ & I \cap (L_1 \cup \dots \cup L_k) = \emptyset \\ 0 & \text{otherwise} \end{cases}$$

A set $\{X_\gamma \mid \gamma \in \Gamma\}$ of operators on a vector space will be defined to be a *bracket set* if for all $\gamma, \gamma' \in \Gamma$, $[X_\gamma, X_{\gamma'}]$ is a multiple of some $X_{\gamma''}$, $\gamma'' \in \Gamma$. If $\{X_\gamma \mid \gamma \in \Gamma\}$ is a bracket set, clearly $\mathfrak{g} = \text{span}\{X_\gamma \mid \gamma \in \Gamma\}$ is a Lie algebra of operators. Our main result is the following:

Theorem 3.1. $\{X_\gamma, Y_\gamma, H_\gamma \mid \gamma \in R^+\}$ is a bracket set inside $GL(V)$ spanning a 14-dimensional Lie algebra $\mathfrak{g} \simeq \mathfrak{g}_2$. The set $\{X_\gamma, Y_\gamma \mid \gamma \in R^+\} \cup \{H_\alpha, H_\beta\}$ is a basis for \mathfrak{g} .

Proof. The proof will require some careful checking involving the G_2 Hexagon. Recall first that the symmetry of the edges and weights of the G_2 Hexagon ensures that for any commutation relation, there is a corresponding relation with all the X operators replaced by Y operators and all the scalar operators H_γ replaced by their negatives. This reduces the number of relations we need check by a factor of two. We proceed in steps.

Step 1. Let us say that $\{\gamma, \gamma'; \gamma''\}$ is a *positive root triple* if all three are positive roots and $\gamma + \gamma' = \gamma''$. The positive root triples are $\{\alpha, \beta; \alpha + \beta\}$, $\{\beta, \alpha + \beta; \alpha + 2\beta\}$, $\{\beta, \alpha + 2\beta; \alpha + 3\beta\}$, $\{\alpha, \alpha + 3\beta; 2\alpha + 3\beta\}$, and $\{\alpha + \beta, \alpha + 2\beta; 2\alpha + 3\beta\}$. For every such triple, there are a number of triangles in the G_2 Hexagon with sides having these directions. Each of these triples leads to six commutation relations of which we need only check three from the above remark. Checking one of these relations involves examining all the triangles associated to that triple and checking consistency of weights between them.

Let us illustrate this for the triple $\{\alpha, \beta; \alpha + \beta\}$. We need check that we can find constants A, B, C such that $[X_\alpha, X_\beta] = AX_{\alpha+\beta}$, $[X_{\alpha+\beta}, Y_\beta] = BX_\alpha$, and $[X_{\alpha+\beta}, Y_\alpha] = CX_\beta$. There are four triangles associated to this triple- they are $\Delta(\phi, \beta, \beta\alpha)$, $\Delta(\beta, \beta\alpha, \beta\alpha\beta)$, $\Delta(\beta\alpha\beta, \beta\alpha\beta\beta, \beta\alpha\beta\beta\alpha)$, and $\Delta(\beta\alpha\beta\beta, \beta\alpha\beta\beta\alpha, \beta\alpha\beta\beta\alpha\beta)$. For one of the triangles, say the first, we observe that the product of the weights of the α and β edges is the weight of the $\alpha + \beta$ edge, and the latter edge is the vector sum of *first* the β edge and *second* the α edge, yielding $[X_\alpha, X_\beta] = X_\alpha X_\beta - X_\beta X_\alpha = -X_{\alpha+\beta}$ when applied to v_ϕ . Now we check that in the other three triangles, exactly the same commutation relation holds. Then we check the same works for the other two relations associated to this triple. Finally we check that the same procedure holds for all positive root triples.

Step 2. There are some potential commutation relations corresponding to the three rectangles in the G_2 Hexagon. These relations involve pairs of roots which are at right angles to each other, namely $\{\beta, 2\alpha + 3\beta\}$, $\{\alpha + \beta, \alpha + 3\beta\}$, $\{\alpha, \alpha + 2\beta\}$ and pairs obtained from them by negating one or both of the entries. Checking that the commutation relations between any X or Y operators involving such a pair is zero amounts to noting that for any one of the rectangles in the G_2 Hexagon, products of weights along adjacent edges is constant.

Step 3. For each of the six positive roots γ , one checks that $H_\gamma = [X_{-\gamma}, X_\gamma]$ is a scalar operator which multiplies each vertex by a number which is the restriction to the Hexagon of a *linear* function in the plane. In particular, H_α corresponds to the linear function which has value -1 on the vertices v_β and $v_{\beta\alpha\beta\beta}$, value 1 on the vertices $v_{\beta\alpha}$ and $v_{\beta\alpha\beta\beta\alpha}$, and value zero on the rest, while H_β corresponds to the linear function which has value 2 at the vertex $v_{\beta\alpha\beta\beta}$, value 1 on the vertices v_β and $v_{\beta\alpha\beta\beta\alpha\beta}$, value zero on $v_{\beta\alpha\beta}$ etc. In particular these two linear functions are linearly independent and so any of the linear functions H_γ is a linear combination of them. From this it follows that $[H_\gamma, X_\gamma]$ is always a multiple of X_γ for any root γ . Finally the Jacobi relation (valid since we are working with operators) and the results from Step 1 show that $[X_{\gamma'}, H_\gamma] = [X_{\gamma'}, [X_{-\gamma}, X_\gamma]] = -[X_{-\gamma}, [X_\gamma, X_{\gamma'}]] - [X_\gamma, [X_{\gamma'}, X_{-\gamma}]$ is a multiple of $X_{\gamma'}$. For example $[X_\gamma, H_\beta] = H_\beta(\gamma)X_\gamma$, where H_β is the value of the linear function H_β on the vector γ .

This shows that altogether the set of operators $\{X_\gamma, Y_\gamma, H_\gamma \mid \gamma \in R^+\}$ is a bracket set of operators, so in particular spans a finite dimensional Lie algebra. It is easy to see that all the $\{X_\gamma, Y_\gamma \mid \gamma \in R^+\}$ are linearly independent, so we get a 14 dimensional Lie algebra. Clearly the span of H_α, H_β is a Cartan subalgebra and the roots with respect to it are exactly $R(G_2)$, in other words, this Lie algebra is \mathfrak{g}_2 . Here is the list of all non-zero commutation relations.

$$\begin{array}{lll}
[X_\beta, X_\alpha] = X_{\alpha+\beta} & [X_{\alpha+\beta}, X_\beta] = 2X_{\alpha+2\beta} & [X_{\alpha+2\beta}, X_\beta] = 3X_{\alpha+3\beta} \\
[X_{\alpha+3\beta}, X_\alpha] = X_{2\alpha+3\beta} & [X_{\alpha+2\beta}, X_{\alpha+\beta}] = 3X_{2\alpha+3\beta} & \\
[Y_\beta, Y_\alpha] = Y_{\alpha+\beta} & [Y_{\alpha+\beta}, Y_\beta] = 2Y_{\alpha+2\beta} & [Y_{\alpha+2\beta}, Y_\beta] = 3Y_{\alpha+3\beta} \\
[Y_{\alpha+3\beta}, Y_\alpha] = Y_{2\alpha+3\beta} & [Y_{\alpha+2\beta}, Y_{\alpha+\beta}] = 3Y_{2\alpha+3\beta} & \\
[X_\alpha, Y_{\alpha+\beta}] = -Y_\beta & [X_\alpha, Y_{2\alpha+3\beta}] = -Y_{\alpha+3\beta} & [X_\beta, Y_{\alpha+3\beta}] = -Y_{\alpha+2\beta} \\
[X_\beta, Y_{\alpha+2\beta}] = -2Y_{\alpha+\beta} & [X_\beta, Y_{\alpha+\beta}] = 3Y_\alpha & [X_{\alpha+\beta}, Y_\alpha] = X_\beta \\
[X_{\alpha+\beta}, Y_\beta] = -3X_\alpha & [X_{\alpha+\beta}, Y_{\alpha+2\beta}] = -2Y_\beta & [X_{\alpha+2\beta}, Y_\beta] = 2X_{\alpha+\beta} \\
[X_{\alpha+2\beta}, Y_{\alpha+\beta}] = 2X_\beta & [X_{\alpha+2\beta}, Y_{\alpha+3\beta}] = Y_\alpha & [X_{\alpha+2\beta}, Y_{2\alpha+3\beta}] = Y_{\alpha+\beta} \\
[X_{\alpha+3\beta}, Y_{2\alpha+3\beta}] = -Y_\alpha & [X_{2\alpha+3\beta}, Y_\alpha] = X_{\alpha+3\beta} & [X_{2\alpha+3\beta}, Y_{\alpha+\beta}] = -X_{\alpha+2\beta} \\
[X_{2\alpha+3\beta}, Y_{\alpha+2\beta}] = -X_{\alpha+\beta} & [X_{2\alpha+3\beta}, Y_{\alpha+3\beta}] = X_\alpha &
\end{array}$$

$$\begin{array}{ll}
[Y_\beta, X_\beta] = H_\beta & [Y_\alpha, X_\alpha] = H_\alpha \\
[Y_{\alpha+\beta}, X_{\alpha+\beta}] = H_{\alpha+\beta} = 3H_\alpha + H_\beta & [Y_{\alpha+2\beta}, X_{\alpha+2\beta}] = H_{\alpha+2\beta} = 2H_\alpha + 3H_\beta \\
[Y_{\alpha+3\beta}, X_{\alpha+3\beta}] = H_{\alpha+3\beta} = H_\alpha + H_\beta & [Y_{2\alpha+3\beta}, X_{2\alpha+3\beta}] = H_{2\alpha+3\beta} = 2H_\alpha + H_\beta \\
[X_\gamma, H_\beta] = H_\beta(\gamma)X_\gamma & [Y_\gamma, H_\beta] = -H_\beta(\gamma)Y_\gamma \\
[X_\gamma, H_\alpha] = H_\alpha(\gamma)X_\gamma & [Y_\gamma, H_\alpha] = -H_\alpha(\gamma)Y_\gamma.
\end{array}$$

□

If \mathfrak{g} is a finite dimensional Lie algebra, it seems interesting to inquire about the existence of spanning bracket sets inside \mathfrak{g} ; do finite ones necessarily exist for example? If so, what is the

minimum number of elements in a spanning bracket set? This is perhaps a useful invariant for Lie algebras.

REFERENCES

- [1] F. Adams, *Lectures on Exceptional Lie Groups*, Chicago Lectures in Mathematics, University of Chicago Press, 1996.
- [2] E. Cartan, *Thèse*, Paris, 1894. 2nd ed., Vuibert, Paris, 1933.
- [3] K. Eriksson, *Strong convergence and a game of numbers*, European J. Combin. **17** (1996), no.4 379–390.
- [4] W. Fulton and J. Harris, *Representation Theory*, Graduate Texts in Mathematics, Springer-Verlag, New York 1991.
- [5] Harish-Chandra, *On some applications of the universal enveloping algebra of a semisimple Lie algebra*, Trans. Amer. Math. Soc. **70**, (1951), 28–96.
- [6] J. Humphreys, *Introduction to Lie Algebras and Representation Theory*, Graduate Texts in Mathematics, Springer-Verlag, New York 1972.
- [7] N. Jacobson, *Lie Algebras*, J. Wiley & Sons, New York 1962.
- [8] N. Jacobson, *Exceptional Lie Algebras*, Lecture Notes in Pure and Applied Mathematics I, Marcel Dekker, New York, 1971.
- [9] W. Killing, *Die Zusammensetzung der stetigen endlichen Transformationsgruppen I,II,III and IV*, Math. Ann. **31** (1888), **33** (1889), **34** (1889) **36** (1890).
- [10] S. Mozes, *Reflection processes on graphs and Weyl groups*, J. Combinatorial Theory A **53** (1990), 128–142.
- [11] R. Proctor, *Minuscule elements of Weyl groups, the numbers game. and d-complete posets*, J. Alg. **213** (1999), no.1, 272–303.
- [12] H. Samelson, *Notes on Lie algebras*, Universitext, Springer-Verlag, New York 1990.
- [13] J. Tits, *Sur les constants de structure et le théoème d'existence des algèbres de Lie semi-simples*, Publ. Math. I.H.E.S. **31**, (1966), 21–58.
- [14] N. J. Wildberger, *A combinatorial construction for simply-laced Lie algebras*, submitted, 2000.
- [15] N. J. Wildberger, *Two new models for the representations of $sl(3)$* , preprint, 2001.
- [16] N. J. Wildberger, *The Mutation Game, Coxeter Graphs, and Partially Ordered Multisets*, preprint, 2001.
E-mail address: n.wildberger@unsw.edu.au

SCHOOL OF MATHEMATICS UNSW SYDNEY NSW 2052 AUSTRALIA