

We dedicate the book to our teachers:

*Peter Gabriel
Shaoxue Liu
Leonard Scott
Jianpan Wang*

A Double Hall Algebra Approach to Affine Quantum Schur–Weyl Theory

Bangming Deng

Jie Du

Qiang Fu[†]

SCHOOL OF MATHEMATICAL SCIENCES, BEIJING NORMAL UNIVERSITY, BEIJING 100875, CHINA.

E-mail address: dengbm@bnu.edu.cn

SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF NEW SOUTH WALES, SYDNEY 2052, AUSTRALIA.

E-mail address: j.du@unsw.edu.au

DEPARTMENT OF MATHEMATICS, TONGJI UNIVERSITY, SHANGHAI, 200092, CHINA.

E-mail address: q.fu@hotmail.com

[†]Corresponding author.

2000 *Mathematics Subject Classification*. Primary 17B37, 20G05, 20C08;
Secondary 16G20, 20C33

Key words and phrases. affine Hecke algebra, affine quantum Schur algebra, cyclic quiver, Drinfeld double, loop algebra, quantum group, Schur–Weyl duality, Ringel–Hall algebra, simple representation

ABSTRACT. Over its one hundred year history, the theory of Schur–Weyl duality and its quantum analogue have made and continue to make profound influences in several areas of mathematics such as Lie theory, representation theory, invariant theory, combinatorial theory, and so on. Recent new developments include, e. g., walled Brauer algebras and rational Schur algebras, quantum Schur superalgebras, and the integral Schur–Weyl duality for types other than A . This book takes an algebraic approach to the affine quantum Schur–Weyl theory.

The book begins with a study of extended Ringel–Hall algebras associated with the cyclic quiver of n vertices and the Green–Xiao Hopf structure on their Drinfeld’s double—the double Ringel–Hall algebra. This algebra is presented in terms of Chevalley type and central generators and is proved to be isomorphic to the quantum loop algebra of the general linear Lie algebra. The rest of the book investigates the affine quantum Schur–Weyl duality on three levels. This includes

- the affine quantum Schur–Weyl reciprocity,
- the bridging role played by the affine quantum Schur algebra between the quantum loop algebra and the corresponding affine Hecke algebra,
- Morita equivalence of certain representation categories,
- presentation of affine quantum Schur algebras, and
- the realization conjecture for the double Ringel–Hall algebra which is proved to be true in the classical case.

Connections with various existing works by Lusztig, Varagnolo–Vasserot, Schiffmann, Hubery, Chari–Pressley, Frenkel–Mukhin, and others are also discussed throughout the book.

Contents

| | |
|---|-----|
| Introduction | 1 |
| Chapter 1. Preliminaries | 7 |
| 1.1. The loop algebra $\widehat{\mathfrak{gl}}_n$ and some notations | 7 |
| 1.2. Representations of cyclic quivers and Ringel–Hall algebras | 9 |
| 1.3. The quantum loop algebra $\mathbf{U}(\widehat{\mathfrak{sl}}_n)$ | 13 |
| 1.4. Three types of generators and associated monomial bases | 16 |
| 1.5. Hopf structure on extended Ringel–Hall algebras | 20 |
| Chapter 2. Double Ringel–Hall algebras of cyclic quivers | 25 |
| 2.1. Drinfeld doubles and the Hopf algebra $\mathfrak{D}_\Delta(n)$ | 25 |
| 2.2. Schiffmann–Hubery generators | 30 |
| 2.3. Presentation of $\mathfrak{D}_\Delta(n)$ | 33 |
| 2.4. Some integral forms | 37 |
| 2.5. The quantum loop algebra $\mathbf{U}(\widehat{\mathfrak{gl}}_n)$ | 40 |
| 2.6. Semisimple generators and commutator formulas | 45 |
| Chapter 3. Affine quantum Schur algebras and the Schur–Weyl reciprocity | 51 |
| 3.1. Cyclic flags: the geometric definition | 51 |
| 3.2. Affine Hecke algebras of type A : the algebraic definition | 55 |
| 3.3. The tensor space interpretation | 61 |
| 3.4. BLM bases and multiplication formulas | 63 |
| 3.5. The $\mathfrak{D}_\Delta(n)$ - $\mathcal{H}_\Delta(r)$ -bimodule structure on tensor spaces | 64 |
| 3.6. A comparison with the Varagnolo–Vasserot action | 72 |
| 3.7. Triangular decompositions of affine quantum Schur algebras | 77 |
| 3.8. Affine quantum Schur–Weyl duality, I | 83 |
| 3.9. Polynomial identity arising from semisimple generators | 86 |
| 3.10. Appendix | 94 |
| Chapter 4. Representations of affine quantum Schur algebras | 99 |
| 4.1. Affine quantum Schur–Weyl duality, II | 100 |
| 4.2. Chari–Pressley’s category equivalence and classification | 103 |
| 4.3. Classification of simple $\mathcal{S}_\Delta(n, r)_{\mathbb{C}}$ -modules: the upward approach | 108 |
| 4.4. Identification of simple $\mathcal{S}_\Delta(n, r)_{\mathbb{C}}$ -modules: the $n > r$ case | 111 |
| 4.5. Application: the set $\mathcal{S}_r^{(n)}$ | 115 |
| 4.6. Classification of simple $\mathcal{S}_\Delta(n, r)_{\mathbb{C}}$ -modules: the downward approach | 116 |
| 4.7. Classification of simple $U_\Delta(n, r)_{\mathbb{C}}$ -modules | 122 |
| Chapter 5. The presentation and realization problems | 125 |

| | | |
|------------|---|-----|
| 5.1. | McGerty's presentation for $\mathbf{U}_\Delta(n, r)$ | 125 |
| 5.2. | Structure of affine quantum Schur algebras | 128 |
| 5.3. | Presentation of $\mathbf{S}_\Delta(r, r)$ | 133 |
| 5.4. | The realization conjecture | 138 |
| 5.5. | Lusztig's transfer maps on semisimple generators | 141 |
| Chapter 6. | The classical ($v = 1$) case | 147 |
| 6.1. | The universal enveloping algebra $\mathcal{U}(\widehat{\mathfrak{gl}}_n)$ | 147 |
| 6.2. | More multiplication formulas in affine Schur algebras | 152 |
| 6.3. | Proof of Conjecture 5.4.1 at $v = 1$ | 156 |
| 6.4. | Appendix: Proof of Proposition 6.2.3 | 160 |
| | Bibliography | 167 |
| | Index | 171 |

Introduction

Quantum Schur–Weyl theory refers to a three-level duality relation. At Level I, it investigates a certain double centralizer property, *the quantum Schur–Weyl reciprocity*, associated with some bimodules of quantum \mathfrak{gl}_n and the Hecke algebra (of type A)—the tensor spaces of the natural representation of quantum \mathfrak{gl}_n (see [41], [20], [26]). This is the quantum version of the well-known Schur–Weyl reciprocity which was beautifully used in H. Weyl’s influential book [77]. The key ingredient of the reciprocity is a class of important finite dimensional endomorphism algebras, the *quantum Schur algebras* or *q -Schur algebra*, whose classical version was introduced by I. Schur over a hundred years ago (see [69], [70]). At Level II, it establishes a certain *Morita equivalence* between quantum Schur algebras and Hecke algebras. Thus, quantum Schur algebras are used to bridge representations of quantum \mathfrak{gl}_n and Hecke algebras. More precisely, they link polynomial representations of quantum \mathfrak{gl}_n with representations of Hecke algebras via the Morita equivalence. The third level of this duality relation is motivated by two simple questions associated with the structure of (associative) algebras. If an algebra is defined by generators and relations, the *realization problem* is to reconstruct the algebra as a vector space with hopefully explicit multiplication formulas on elements of a basis; while, if an algebra is defined in term of a vector space such as an endomorphism algebra, it is natural to seek their generators and defining relations.

As one of the important problems in quantum group theory, the realization problem is to construct a quantum group in terms of a vector space and certain multiplication rules on basis elements. This problem is crucial to understand their structure and representations (see [45, p. xiii] for a similar problem for Kac–Moody Lie algebras and [58] for a solution in the symmetrizable case). Though the Ringel–Hall algebra realization of the \pm -part of quantum enveloping algebras associated with symmetrizable Cartan matrices is an important breakthrough in the early nineties of the last century, especially for the introduction of the geometric approach to the theory, the same problem for the entire quantum groups is far from completion. However, Beilinson–Lusztig–MacPherson (BLM) [3] solved the problem for quantum \mathfrak{gl}_n by exploring further properties coming from the quantum Schur–Weyl reciprocity. On the other hand, as endomorphism algebras and as homomorphic images of quantum \mathfrak{gl}_n , it is natural to look for presentations for quantum Schur algebras via the presentation of quantum \mathfrak{gl}_n . This problem was first considered in [17] (see also [25]). Thus, as a particular feature in the type A theory, realizing quantum \mathfrak{gl}_n and presenting quantum Schur algebras form the Level III of this duality relation. For a complete account of the quantum Schur–Weyl theory and further references, see Parts 3 and 5 of [11] (see also [16] for more applications).

There are several developments in the establishment of an affine analogue of the quantum Schur–Weyl theory. Soon after BLM’s work, Ginzburg and Vasserot [31, 75] used a geometric and K -theoretic approach to investigate *affine quantum Schur algebras*¹ as homomorphic images of *quantum loop algebra* $\mathbf{U}(\widehat{\mathfrak{gl}}_n)$ of \mathfrak{gl}_n in the sense of Drinfeld’s new presentation [19], called quantum affine \mathfrak{gl}_n (at level 0) in this book. This establishes at Level I the first centralizer property for the affine analogue of the quantum Schur–Weyl reciprocity. Six years later, investigations around affine quantum Schur algebras focused on their different definitions and, hence, different applications. For example, Lusztig [54] generalized the fundamental multiplication formulas [3, 3.4] for quantum Schur algebras to the affine case and showed that the “extended” quantum affine \mathfrak{sl}_n , $\mathbf{U}_\Delta(n)$, does not map onto affine quantum Schur algebras; Varagnolo–Vasserot [73] investigated Ringel–Hall algebra actions on tensor spaces and described the geometrically-defined affine quantum Schur algebras in terms of the endomorphism algebras of tensor spaces. Moreover, they proved that the tensor space definition coincides with Green’s definition [34] via q -permutation modules. Some progress on the second centralizer property has also been made recently by Pouchin [59]. The approaches used in these works are mainly geometric. However, like the non-affine case, there would be more favorable algebraic and combinatorial approaches.

At Level II, representations at non-roots-of-unity of quantum affine \mathfrak{sl}_n and \mathfrak{gl}_n over the complex number field \mathbb{C} , including classifications of finite dimensional simple modules, have been thoroughly investigated by Chari–Pressley [5, 6, 7], and Frenkel–Mukhin [27] in terms of Drinfeld polynomials. Moreover, an equivalence between the module category of the Hecke algebra $\mathcal{H}_\Delta(r)_\mathbb{C}$ and a certain full subcategory of quantum affine \mathfrak{sl}_n (resp., \mathfrak{gl}_n) has also been established algebraically by Chari–Pressley [8] (resp., geometrically by Ginzburg–Reshetikhin–Vasserot [30]) under the condition $n > r$ (resp., $n \geq r$). Note that the approach in [30] uses intersection cohomology complexes. It would be interesting to know how affine quantum Schur algebras would play a role in these works.

Much less progress has been made at Level III. When $n > r$, Doty–Green [17] and McGerty [56] have found a presentation for affine quantum Schur algebras, while the last two authors of the book have investigated the realization problem in [23], where they first developed an approach without using the stabilization property, a key property used in the BLM approach, and presented an ideal candidate for the realization of quantum affine \mathfrak{gl}_n .

This book attempts to establish the affine quantum Schur–Weyl theory as a whole and is an outcome of *algebraically* understanding the works mentioned above.

First, building on Schiffmann [67] and Hubery [38], our starting point is to present the double Ringel–Hall algebra $\mathfrak{D}_\Delta(n)$ of the cyclic quiver with n vertices in terms of Chevalley type generators together with infinitely many central generators. Thus, we obtain a central subalgebra $\mathbf{Z}_\Delta(n)$ such that $\mathfrak{D}_\Delta(n) = \mathbf{U}_\Delta(n)\mathbf{Z}_\Delta(n) \cong \mathbf{U}_\Delta(n) \otimes \mathbf{Z}_\Delta(n)$. We then establish an isomorphism between $\mathfrak{D}_\Delta(n)$ and Drinfeld’s quantum affine \mathfrak{gl}_n

¹Perhaps, it should be called quantum affine Schur algebras. Since our purpose is to establish an affine analogue of the quantum Schur–Weyl theory, this terminology seems more appropriate to reflect this.

in the sense of [19]. In this way, we easily obtain an action on the tensor space which upon restriction coincides with the Ringel–Hall algebra action defined geometrically by Varagnolo–Vasserot [73] and commutes with the affine Hecke algebra action.

Second, by a thorough investigation of a BLM type basis for affine quantum Schur algebras, we introduce certain triangular relations for the corresponding structure constants and, hence, a triangular decomposition for affine quantum Schur algebras. With this decomposition, we establish explicit algebra epimorphisms $\xi_r = \xi_{r, \mathbb{Q}(v)}$ from the double Ringel–Hall algebra $\mathfrak{D}_\Delta(n)$ to affine quantum Schur algebras $\mathcal{S}_\Delta(n, r) := \mathcal{S}_\Delta(n, r)_{\mathbb{Q}(v)}$ for all $r \geq 0$. This algebraic construction has several nice applications, especially at Levels II and III. For example, the homomorphic image of commutator formulas for semisimple generators gives rise to a beautiful polynomial identity whose combinatorial proof remains mysterious.

Like the quantum Schur algebra case, we will establish for $n \geq r$ a Morita equivalence between affine quantum Schur algebras $\mathcal{S}_\Delta(n, r)_{\mathbb{F}}$ and affine Hecke algebras $\mathcal{H}_\Delta(r)_{\mathbb{F}}$ of type A over a field \mathbb{F} with a non-root-of-unity parameter. As a by-product, we prove that every simple $\mathcal{S}_\Delta(n, r)_{\mathbb{F}}$ -module is finite dimensional. Thus, applying the classification of simple $\mathcal{H}_\Delta(r)_{\mathbb{C}}$ -modules by Zelevinsky [81] and Rogawski [66] yields a classification of simple $\mathcal{S}_\Delta(n, r)_{\mathbb{C}}$ -modules. Hence, inflation via the epimorphisms $\xi'_{r, \mathbb{C}}$ gives many finite dimensional simple $U_{\mathbb{C}}(\widehat{\mathfrak{gl}}_n)$ -modules. We will also use $\xi'_{r, \mathbb{C}}$ together with the action on tensor spaces and a result of Chari–Pressley to prove that finite dimensional simple polynomial representations of $U_{\mathbb{C}}(\widehat{\mathfrak{gl}}_n)$ are all inflations of simple $\mathcal{S}_\Delta(n, r)_{\mathbb{C}}$ -modules. In this way, we can see the bridging role played by affine quantum Schur algebras between representations of quantum affine \mathfrak{gl}_n and those of affine Hecke algebras. Moreover, we obtain a classification of simple $\mathcal{S}_\Delta(n, r)_{\mathbb{C}}$ -modules in terms of Drinfeld polynomials and, when $n > r$, we identify them with those arising from simple $\mathcal{H}_\Delta(r)_{\mathbb{C}}$ -modules.

Our findings also show that, if we regard the category $\mathcal{S}_\Delta(n, r)_{\mathbb{C}}\text{-Mod}$ of $\mathcal{S}_\Delta(n, r)_{\mathbb{C}}$ -modules as a full subcategory of $U_{\mathbb{C}}(\widehat{\mathfrak{gl}}_n)$ -modules, this category is quite different from the category $\mathcal{C}^{hi} \cap \mathcal{C}'$ considered in [52, §6.2]. For example, the latter is completely reducible and simple objects are usually infinite dimensional, while $\mathcal{S}_\Delta(n, r)_{\mathbb{C}}\text{-Mod}$ is not completely reducible and all simple objects are finite dimensional. As observed in [22, Rem. 9.4(2)] for quantum \mathfrak{gl}_∞ and infinite quantum Schur algebras, this is another kind of phenomenon of infinite type in contrast with the finite type case.

The discussion of the realization and presentation problems is also based on the algebra epimorphisms ξ_r and relies on the use of semisimple generators and indecomposable generators for $\mathfrak{D}_\Delta(n)$ which are crucial to understand the integral structure and multiplication formulas. We first use the new presentation for $\mathfrak{D}_\Delta(n)$ to give a decomposition for $\mathcal{S}_\Delta(n, r) = \mathbf{U}_\Delta(n, r)\mathbf{Z}_\Delta(n, r)$ into a product of two subalgebras, where $\mathbf{Z}_\Delta(n, r)$ is a central subalgebra and $\mathbf{U}_\Delta(n, r)$ is the homomorphic image of $\mathbf{U}_\Delta(n)$, the extended quantum affine \mathfrak{sl}_n . By a close look at this structure, we manage to get a presentation for $\mathcal{S}_\Delta(n, r)$ for all $r \geq 1$ and acknowledge that the presentation problem is very complicated in the $n < r$ case. On the other hand, we formulate a realization conjecture suggested by the work [23] and prove the conjecture in the classical ($v = 1$) case.

We remark that, unlike the geometric approach in which the ground ring must be a field or mostly the complex number field \mathbb{C} , the algebraic, or rather, the representation-theoretic approach we used in the book works largely over a ring or mostly the integral Laurent polynomial ring $\mathbb{Z}[v, v^{-1}]$.

We organize the book as follows.

In the first preliminary chapter, we introduce in §1.4 three different types of generators and their associated monomial bases for the Ringel–Hall algebras of cyclic quivers, and display in §1.5 the Green–Xiao Hopf structure on the extended version of these algebras.

Chapter 2 introduces a new presentation using Chevalley generators for Drinfeld’s quantum loop algebra $\mathbf{U}(\widehat{\mathfrak{gl}}_n)$ of \mathfrak{gl}_n . This is achieved by constructing the presentation for the double Ringel–Hall algebra $\mathfrak{D}_\Delta(n)$ associated with cyclic quivers (Theorem 2.3.1), based on works of Schiffmann and Hubery, and by lifting Beck’s algebra monomorphism from the quantum $\widehat{\mathfrak{sl}}_n$ with a Drinfeld–Jimbo presentation into $\mathbf{U}(\widehat{\mathfrak{gl}}_n)$ to obtain an isomorphism between $\mathfrak{D}_\Delta(n)$ and $\mathbf{U}(\widehat{\mathfrak{gl}}_n)$ (Theorem 2.5.3).

Chapter 3 investigates the structure of affine quantum Schur algebras. We first recall the geometric definition by Ginzburg–Vasserot and Lusztig, the Hecke algebra definition by R. Green, and the tensor space definition by Varagnolo–Vasserot. Using the Chevalley generators of $\mathfrak{D}_\Delta(n)$, we easily obtain an action on the $\mathbb{Q}(v)$ -space Ω with a basis indexed by \mathbb{Z} and, hence, an action of $\mathfrak{D}_\Delta(n)$ on $\Omega^{\otimes r}$ (§3.5). We prove that this action commutes with the affine Hecke algebra action defined in [73]. Moreover, we show that the restriction of the action to the \pm -part of $\mathfrak{D}_\Delta(n)$ (i. e., to the corresponding Ringel–Hall algebra) coincides with the Ringel–Hall algebra action geometrically defined by Varagnolo–Vasserot (Theorem 3.6.3). As an application of this coincidence, the commutator formula associated with semisimple generators, arising from the skew-Hopf pairing, gives rise to certain polynomial identity associated with a pair of elements $\lambda, \mu \in \mathbb{N}_\Delta^n$ (Corollary 3.9.6). The main result of the chapter is an elementary proof of the surjective homomorphism ξ_r from the double Ringel–Hall algebra $\mathfrak{D}_\Delta(n)$, i. e., the quantum loop algebra $\mathbf{U}(\widehat{\mathfrak{gl}}_n)$, onto the affine quantum Schur algebra $\mathcal{S}_\Delta(n, r)$ (Theorem 3.8.1). The approach we used is the establishment of a triangular decomposition of $\mathcal{S}_\Delta(n, r)$ (Theorem 3.7.7) through an analysis of the BLM type bases.

In Chapter 4, we discuss the representation theory of affine quantum Schur algebras over \mathbb{C} and its connection to polynomial representations of quantum affine \mathfrak{gl}_n and representations of affine Hecke algebras. We first establish a category equivalence between the module categories $\mathcal{S}_\Delta(n, r)_\mathbb{C}\text{-Mod}$ and $\mathcal{H}_\Delta(r)_\mathbb{C}\text{-Mod}$ for $n \geq r$ (Theorem 4.1.3). As an application, we will reinterpret Chari–Pressley’s category equivalence ([8, Th. 4.2]) between (level r) representations of $U_\mathbb{C}(\widehat{\mathfrak{sl}}_n)$ and those of affine Hecke algebras $\mathcal{H}_\Delta(r)_\mathbb{C}$, where $n > r$, in terms of representations of $\mathcal{S}_\Delta(n, r)_\mathbb{C}$ (Proposition 4.2.1). We then develop two approaches to the classification of simple $\mathcal{S}_\Delta(n, r)_\mathbb{C}$ -modules. In the so-called upward approach, we use the classification of simple $\mathcal{H}_\Delta(r)_\mathbb{C}$ -modules of Zelevinsky and Rogawski to classify simple $\mathcal{S}_\Delta(n, r)_\mathbb{C}$ -modules (Theorems 4.3.4 and 4.5.3), while in the downward approach, we determine the classification of simple $\mathcal{S}_\Delta(n, r)_\mathbb{C}$ -modules (Theorem 4.6.8) in terms of simple polynomial representations of $U_\mathbb{C}(\widehat{\mathfrak{gl}}_n)$. When $n > r$, we prove an identification theorem (Theorem 4.4.2)

for the two classifications. Finally, in §4.7, a classification of finite dimensional simple $U_\Delta(n, r)_\mathbb{C}$ -modules is also completed and its connections to finite dimensional simple $U_\mathbb{C}(\widehat{\mathfrak{sl}}_n)$ -modules and finite dimensional simple (polynomial) $U_\mathbb{C}(\widehat{\mathfrak{gl}}_n)$ -modules are also discussed.

We move on to look at the presentation and realization problems in Chapter 5. We first observe $\mathcal{S}_\Delta(n, r) = \mathbf{U}_\Delta(n, r)\mathbf{Z}_\Delta(n, r)$, where $\mathbf{U}_\Delta(n, r)$ and $\mathbf{Z}_\Delta(n, r)$ are homomorphic images of $\mathbf{U}_\Delta(n)$ and $\mathbf{Z}_\Delta(n)$, respectively, and that $\mathbf{Z}_\Delta(n, r) \subseteq \mathbf{U}_\Delta(n, r)$ if and only if $n > r$. A presentation for $\mathbf{U}_\Delta(n, r)$ is given in [56] (see also [18] for the $n > r$ case). Building on McGerty's presentation, we first give a Drinfeld–Jimbo type presentation for the subalgebra $\mathbf{U}_\Delta(n, r)$ (Theorem 5.1.3). We then describe a presentation for the central subalgebra $\mathbf{Z}_\Delta(n, r)$ as a Laurent polynomial ring in one indeterminate over a polynomial ring in $r - 1$ indeterminates over $\mathbb{Q}(v)$. We manage to describe a presentation for $\mathcal{S}_\Delta(n, r)$ for all $r \geq 1$ (Theorem 5.3.5) by adding an extra generator (and its inverse) together with an additional set of relations on top of the relations given in Theorem 5.1.3. What we will see from this case is that the presentation for $\mathcal{S}_\Delta(n, r)$ with $r > n$ can be very complicated.

We discuss the realization problem from §5.4 onwards. We first describe the modified BLM approach developed in [23]. With some supporting evidence, we then formulate the realization conjecture (Conjecture 5.4.2) as suggested in [23, 5.5(2)], and state its classical ($v = 1$) version. We end the chapter with a closer look at Lusztig's transfer maps [55] by displaying some explicit formulas for their action on the semisimple generators for $\mathcal{S}_\Delta(n, r)$ (Corollary 5.5.2). These formulas show also that the homomorphism from $\mathbf{U}(\widehat{\mathfrak{sl}}_n)$ to $\varprojlim \mathcal{S}_\Delta(n, n + m)$ induced by the transfer maps cannot either be extended to the double Ringel–Hall algebra $\mathcal{D}_\Delta(n)$. (Lusztig already pointed out that it cannot be extended to $\mathbf{U}_\Delta(n)$.) This somewhat justifies why a direct product is used in the realization conjecture.

In the last Chapter 6, we prove the realization conjecture for the classical ($v = 1$) case. The key step to the proof is the establishment of more multiplication formulas (Proposition 6.2.3) between homogeneous indecomposable generators and an arbitrary BLM type basis element. As a by-product, we display a basis for the universal enveloping algebra of the loop algebra of \mathfrak{gl}_n (Theorem 6.3.4) together with explicit multiplication formulas between generators and arbitrary basis elements (Corollary 6.3.5).

There are two Appendices §§3.10, 6.4 which collect a number of lengthy calculations used in some proofs.

Conjectures and problems. There are quite a few conjectures and problems throughout the book. The conjectures are mostly natural generalizations to the affine case, for example, the realization conjecture 5.4.2 and the conjectures in §3.8 on an integral form for double Ringel–Hall algebras and the second centralizer property in the affine quantum Schur–Weyl reciprocity. Some problems are designed to seek further solutions to certain questions such as “quantum Serre relations” for semisimple generators (Problem 2.6.4), the Affine Branching Rule (Problem 4.3.6), and further identification of simple modules from different classifications (Problem 4.6.11). There

are also problems for seeking different proofs. Problems 3.4.3 and 6.4.2 form a key step towards the proof of the realization conjecture.

Notational scheme. For most of the notations throughout the book, if it involves a subscript Δ or a superscript Δ , it indicates that the same notation without Δ has been used in the non-affine case, say, in [3], [11], [32], etc. Here the triangle Δ depicts the cyclic Dynkin diagram of affine type A .

For a ground ring \mathcal{Z} and a \mathcal{Z} -module (or a \mathcal{Z} -algebra) \mathcal{A} , we often use the notation $\mathcal{A}_{\mathbb{F}} := \mathcal{A} \otimes \mathbb{F}$ to represent the object obtained by *base change* to a field \mathbb{F} , which itself is a \mathcal{Z} -module. In particular, if $\mathcal{Z} = \mathbb{Z}[v, v^{-1}]$, then we write \mathcal{A} for $\mathcal{A}_{\mathbb{Q}(v)}$.

Acknowledgements. The main results of the book have been presented by the authors at the following conferences and workshops. We would like to thank the organizers for the opportunities of presenting our work.

- Conference on Perspectives in Representation Theory, Cologne, September 2009;
- International Workshop on Combinatorial and Geometric Approach to Representation Theory, Seoul National University, September 2009;
- 2010 ICM Satellite Conference, Bangalore, August 2010;
- 12th National Algebra Conference, Lanzhou, June 2010;
- Southeastern Lie Theory Workshop: Finite and algebraic groups and Leonard Scott Day, Charlottesville, June 2011;
- 55th Annual Meeting of the Australian Mathematical Society, Wollongong, September 2011.

The research was partially supported by the Australian Research Council, the Natural Science Foundation of China, the 111 Program of China, the Program NCET, the Fundamental Research Funds for the Central Universities of China, and the UNSW Goldstar Award. The first three and last two chapters were written while Deng and Fu were visiting the University of New South Wales at the various times. The hospitality and support of UNSW are gratefully acknowledged.

The second author would like to thank Alexander Kleshchev and Arun Ram for helpful comments on the Affine Branching Rule (4.3.5.1), and Vyjayanthi Chari for several discussions and explanations on the paper [8] and some related topics. He would also like to thank East China Normal University, Universities of Mainz, Virginia and Auckland for their hospitality during his sabbatical leave in the second half of 2009.

Finally, for all their help, encouragement, and infinite patience, we thank our wives and children: Wenlian Guo and Zhuoran Deng; Chunli Yu, Andy Du, and Jason Du; Shanshan Xia.

Bangming Deng
Jie Du
Qiang Fu

Sydney
5 December 2011