

# A CONNECTION BETWEEN $\pi$ AND $\phi$

MICHAEL D. HIRSCHHORN

ABSTRACT. We find an expression for  $\pi$  as a limit involving the golden ratio,  $\phi$ .

## 1. INTRODUCTION

We prove that

$$\sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{n} \sim \frac{\phi^{5n+\frac{5}{2}}}{2\pi n^4 \sqrt{5}} \left( 1 - \frac{5-\sqrt{5}}{10n} + \frac{13-5\sqrt{5}}{50n^2} - \frac{175-83\sqrt{5}}{1250n^3} + \frac{437-205\sqrt{5}}{6250n^4} - \dots \right)$$

as  $n \rightarrow \infty$ , where  $\phi$  is the golden ratio, and consequently,

$$\frac{1}{\pi} = \lim_{n \rightarrow \infty} 2n^4 \sqrt{5} \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{n} / \phi^{5n+\frac{5}{2}}.$$

In order to carry out this program, we use the methods developed in two earlier papers on the Apéry numbers [2], [3].

## 2. THE DOMINANT TERM

The first step is to find the value of  $k$  for which the term  $\binom{n}{k}^2 \binom{n+k}{n}$  is a maximum. We do this by setting

$$\binom{n}{k}^2 \binom{n+k}{n} = \binom{n}{k+1}^2 \binom{n+k+1}{n}.$$

This yields

$$\frac{n!(n+k)!}{k!^3(n-k)!^2} = \frac{n!(n+k+1)!}{(k+1)!^3(n-k-1)!^2},$$

or,

$$(k+1)^3 = (n+k+1)(n-k)^2.$$

If we suppose  $k = \theta n$ , where  $\theta$  is to be determined, and divide by  $n^3$ , we find

$$\left(\theta + \frac{1}{n}\right)^3 = \left(1 + \theta + \frac{1}{n}\right)(1 - \theta)^2.$$

If we let  $n \rightarrow \infty$ , this becomes

$$\theta^3 = (1 + \theta)(1 - \theta)^2,$$

or,

$$1 - \theta - \theta^2 = 0.$$

It follows that

$$\theta = \frac{\sqrt{5} - 1}{2} = \frac{1}{\phi},$$

where  $\phi$  is the golden ratio.

Thus, the value of  $k$  that we seek is given by

$$k \approx \theta n$$

where  $\theta = \frac{1}{\phi}$ .

At  $k \approx \theta n$ , the value of the term is

$$\begin{aligned} H &= \binom{n}{\theta n}^2 \binom{n + \theta n}{n} \\ &= \frac{n!(n + \theta n)!}{(\theta n)!^3 (n - \theta n)!^2} \\ &\approx \frac{\sqrt{2\pi n} \left(\frac{n}{e}\right)^n \sqrt{2\pi(1 + \theta)n} \left(\frac{(1 + \theta)n}{e}\right)^{(1 + \theta)n}}{\left(\sqrt{2\pi\theta n} \left(\frac{\theta n}{e}\right)^{\theta n}\right)^3 \left(\sqrt{2\pi(1 - \theta)n} \left(\frac{(1 - \theta)n}{e}\right)^{(1 - \theta)n}\right)^2} \\ &= \left(\frac{1}{\sqrt{2\pi n}}\right)^3 \frac{\sqrt{1 + \theta}}{(\sqrt{\theta})^3 (\sqrt{1 - \theta})^2} \left(\frac{(1 + \theta)^{1 + \theta}}{\theta^{3\theta} (1 - \theta)^{2(1 - \theta)}}\right)^n \end{aligned}$$

after considerable simplification.

Now,

$$1 - \theta = \theta^2 \quad \text{and} \quad 1 + \theta = \frac{1}{\theta}$$

so

$$\frac{\sqrt{1 + \theta}}{(\sqrt{\theta})^3 (\sqrt{1 - \theta})^2} = \frac{1}{1 - \theta} \sqrt{\frac{1 + \theta}{\theta^3}} = \frac{1}{\theta^2} \sqrt{\frac{1}{\theta^4}} = \frac{1}{\theta^4} = \phi^4$$

and

$$\frac{(1 + \theta)^{1 + \theta}}{\theta^{3\theta} (1 - \theta)^{2(1 - \theta)}} = \frac{1}{\theta^{1 + \theta} \theta^{3\theta} (\theta^2)^{2(1 - \theta)}} = \frac{1}{\theta^5} = \phi^5.$$

So

$$H \approx \frac{\phi^{5n+4}}{(2\pi n)^{\frac{3}{2}}}.$$

(See fig. 1.)

At points near  $\theta n$ , the terms of the sum are given by

$$\begin{aligned} \binom{n}{\theta n + k}^2 \binom{n + \theta n + k}{n} &= H \cdot \binom{n}{\theta n + k}^2 \binom{n + \theta n + k}{n} / \binom{n}{\theta n}^2 \binom{n + \theta n}{n} \\ &= H \cdot \frac{n!(n + \theta n + k)!}{(\theta n + k)!^3 (n - \theta n - k)!^2} / \frac{n!(n + \theta n)!}{(\theta n)!^3 (n - \theta n)!^2} \\ &= H \cdot \frac{(n + \theta n + 1) \cdots (n + \theta n + k)(n - \theta n)^2 \cdots (n - \theta n - k + 1)^2}{(\theta n + 1)^3 (\theta n + 2)^3 \cdots (\theta n + k)^3} \end{aligned}$$




FIGURE 1. The case  $n = 1000$ , showing the points  $(k, \binom{n}{k}^2 \binom{n+k}{n})$  for  $580 \leq k \leq 660$ , together with the vertical  $x = \frac{n}{\phi}$ ,  $0 \leq y \leq \frac{\phi^{5n+4}}{(2\pi n)^{\frac{3}{2}}}$

$$\begin{aligned}
&= H \cdot \frac{\left(1 + \frac{1}{(1+\theta)n}\right) \cdots \left(1 + \frac{k}{(1+\theta)n}\right) \left(1 - \frac{1}{(1-\theta)n}\right)^2 \cdots \left(1 - \frac{k-1}{(1-\theta)n}\right)^2}{\left(1 + \frac{1}{\theta n}\right)^3 \cdots \left(1 + \frac{k}{\theta n}\right)^3} \\
&\approx H \exp \left\{ \frac{1}{(1+\theta)n} (1 + \cdots + k) - \frac{2}{(1-\theta)n} (1 + \cdots + (k-1)) - \frac{3}{\theta n} (1 + \cdots + k) \right\} \\
&\approx H \exp \left\{ -\frac{k^2}{2n} \left( \frac{2}{1-\theta} + \frac{3}{\theta} - \frac{1}{1+\theta} \right) \right\} \\
&= H \exp \left\{ -\frac{k^2}{2n} \phi^3 \sqrt{5} \right\}
\end{aligned}$$

since

$$\frac{(1+\theta)^k (1-\theta)^{2k}}{\theta^{3k}} = \frac{(\theta^2)^{2k}}{\theta^k \cdot \theta^{3k}} = 1$$

and

$$\begin{aligned}
\frac{2}{1-\theta} + \frac{3}{\theta} - \frac{1}{1+\theta} &= \frac{2}{\theta^2} + \frac{3}{\theta} - \theta = \frac{2+3\theta-\theta^3}{\theta^2} = \frac{2+3\theta-\theta(1-\theta)}{\theta^2} \\
&= \frac{2+2\theta+\theta^2}{\theta^2} = (2+2\theta+\theta^2)\phi^2 = 2\phi^2 + 2\phi + 1 = 4\phi + 3 \\
&= 4 \left( \frac{\sqrt{5}+1}{2} \right) + 3 = 2\sqrt{5} + 5 = (2+\sqrt{5})\sqrt{5} = \phi^3 \sqrt{5}.
\end{aligned}$$

Thus, the terms are essentially distributed normally, with  $\sigma^2 = \frac{n}{\phi^3 \sqrt{5}}$ , and the sum is given by

$$\sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{n} \approx H \int_{-\infty}^{\infty} \exp \left\{ -\frac{x^2}{2\sigma^2} \right\} dx = H \cdot \sigma \sqrt{2\pi}$$

$$\begin{aligned} &\approx \frac{\phi^{5n+4}}{(2\pi n)^{\frac{3}{2}}} \cdot \frac{\sqrt{n}}{\phi^{\frac{3}{2}} \sqrt[4]{5}} \sqrt{2\pi} \\ &= \frac{\phi^{5n+\frac{5}{2}}}{2\pi n \sqrt[4]{5}}, \end{aligned}$$

as claimed. (See fig. 2.)




FIGURE 2. The case  $n = 1000$ , showing the points  $(k, \binom{n}{k}^2 \binom{n+k}{n})$  for  $550 \leq k \leq 690$ , together with the approximating normal,  $y = \frac{\phi^{5n+4}}{(2\pi n)^{\frac{3}{2}}} \exp \left\{ -\frac{\phi^3 \sqrt{5}}{2n} \left( x - \frac{n}{\phi} \right)^2 \right\}$ .

### 3. THE CORRECTION TERM

Let  $s_n = \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{n}$ . It was stated by Apéry [1] and proved by A. van der Poorten (see §4) that  $s_n$  satisfies the recurrence

$$(n+1)^2 s_{n+1} - (11n^2 + 11n + 3)s_n - n^2 s_{n-1} = 0,$$

or,

$$\left(1 + \frac{1}{n}\right)^2 s_{n+1} - \left(11 + \frac{11}{n} + \frac{3}{n^2}\right) s_n - s_{n-1} = 0.$$

We now suppose that

$$s_n = Cn^{-1} \Phi^n \left(1 + \frac{a_1}{n} + \frac{a_2}{n^2} + \frac{a_3}{n^3} + \dots\right),$$

where  $C = \frac{\phi^{\frac{5}{2}}}{2\pi \sqrt[4]{5}}$  and  $\Phi = \phi^5$ , and substitute into the recurrence, to obtain

$$\left(1 + \frac{1}{n}\right)^2 C(n+1)^{-1} \Phi^{n+1} \left(1 + \frac{a_1}{n+1} + \frac{a_2}{(n+1)^2} + \frac{a_3}{(n+1)^3} + \dots\right)$$

$$\begin{aligned}
& - \left( 11 + \frac{11}{n} + \frac{3}{n^2} \right) C n^{-1} \Phi^n \left( 1 + \frac{a_1}{n} + \frac{a_2}{n^2} + \frac{a_3}{n^3} + \dots \right) \\
& - C(n-1)^{-1} \Phi^{n-1} \left( 1 + \frac{a_1}{n-1} + \frac{a_2}{(n-1)^2} + \frac{a_3}{(n-1)^3} + \dots \right) = 0.
\end{aligned}$$

If we now divide by  $C\Phi^n$ , and multiply by  $n$ , we find

$$\begin{aligned}
& \left( 1 + \frac{1}{n} \right) \Phi \left( 1 + \frac{a_1}{n+1} + \frac{a_2}{(n+1)^2} + \frac{a_3}{(n+1)^3} + \dots \right) \\
& - \left( 11 + \frac{11}{n} + \frac{3}{n^2} \right) \left( 1 + \frac{a_1}{n} + \frac{a_2}{n^2} + \frac{a_3}{n^3} + \dots \right) \\
& - \left( 1 - \frac{1}{n} \right)^{-1} \Phi^{-1} \left( 1 + \frac{a_1}{n-1} + \frac{a_2}{(n-1)^2} + \frac{a_3}{(n-1)^3} + \dots \right) = 0.
\end{aligned}$$

If we set  $\frac{1}{n} = u$ ,  $\frac{1}{n+1} = \frac{u}{1+u}$ ,  $\frac{1}{n-1} = \frac{u}{1-u}$ , and expand in powers of  $u$ , we find

$$\begin{aligned}
& \Phi(1+u)(1+a_1u+(a_2-a_1)u^2+(a_3-2a_2+a_1)u^3+\dots) \\
& - (11+11u+3u^2)(1+a_1u+a_2u^2+a_3u^3+\dots) \\
& - \Phi^{-1}(1+u+u^2+u^3+\dots)(1+a_1u+(a_2+a_1)u^2+(a_3+2a_2+a_1)u^3+\dots) \\
& = 0.
\end{aligned}$$

We now set the coefficients of the powers of  $u$  equal to zero, and solve for  $a_1$ ,  $a_2$ ,  $a_3$  and so on. The constant term and the coefficient of  $u$  are automatically zero, because we had  $\Phi$  correct and the factor  $n^{-1}$  correct. The coefficient of  $u^2$  is

$$\Phi a_2 - (11a_2 + 11a_1 + 3) - \Phi^{-1}(a_2 + 2a_1 + 1) = 0,$$

or,

$$-(11 + 2\Phi^{-1})a_1 - (3 + \Phi^{-1}) = 0.$$

We find

$$a_1 = -\frac{3 + \Phi^{-1}}{11 + 2\Phi^{-1}} = -\frac{3 + \left(\frac{5\sqrt{5} - 11}{2}\right)}{11 + 2\left(\frac{5\sqrt{5} - 11}{2}\right)} = -\frac{5\sqrt{5} - 5}{10\sqrt{5}} = -\frac{5 - \sqrt{5}}{10}.$$

If we continue in the same way, we find

$$a_2 = \frac{13 - 5\sqrt{5}}{50}, \quad a_3 = -\frac{175 - 83\sqrt{5}}{1250}, \quad a_4 = \frac{437 - 205\sqrt{5}}{6250},$$

and so on.

This completes the proof.

## 4. THE RECURRENCE

A. van der Poorten's proof [5] goes as follows.

If we define

$$f(k) = (k^2 + (6n + 3)k - (11n^2 + 9n + 2)) \binom{n}{k}^2 \binom{n+k}{n}$$

and

$$g(n) = \binom{n}{k}^2 \binom{n+k}{n}$$

then it is easy to verify that

$$f(k) - f(k-1) = (n+1)^2 g(n+1) - (11n^2 + 11n + 3)g(n) - n^2 g(n-1).$$

The recurrence follows on summing over  $k$  from 0 to  $n+1$ .

Following the work of Sister Celine Fasenmyer and Petrovsek, Wilf and Zeilberger [4], the discovery of such identities is routine.

## REFERENCES

- [1] R. Apéry, Irrationalité de  $\zeta(2)$ ,  $\zeta(3)$ , *Astérisque* 61(1979), 11–13.
- [2] M. D. Hirschhorn, Estimating the Apéry numbers, *Fibonacci Quarterly* 50(2)(2012), 129–131.
- [3] M. D. Hirschhorn, Estimating the Apéry numbers II, *Fibonacci Quarterly*, to appear.
- [4] M. Petrovsek, H. Wilf and D. Zeilberger, *A=B*, A. K. Peters Ltd., Wellesley MA, 1996.
- [5] A. van der Poorten, A proof that Euler missed, *Mathematical Intelligencer* 1(1979), 195–203.

AMS Classification Numbers: 41A60

SCHOOL OF MATHEMATICS AND STATISTICS, UNSW, SYDNEY, AUSTRALIA 2052